


First Name: _____

Last Name: _____

LO3 - EQ - Mass Spectrometers and Millikan's Experiment - ANSWER KEY

Use the following information to answer Q1:



Q1: What is the most probable location for data collected in a Millikan-style experiment?

- a. Location 1
- b. Location 2
- c. Location 3
- d. Location 4

Linear relationship.

$$mg = q|\vec{E}| \quad \text{or} \quad q = \frac{g}{|\vec{E}|}m + 0 \quad (\text{in } y=mx+b \text{ format})$$

So no y-intercept. Also since charge is quantized, the only options are ones that are equal to ne , where n is an integer and e is the elementary charge (1.60×10^{-19} C).

Use the following information to answer Q2:

Mass Spectrometer Equation Configurations			
1 -	$q\Delta V = \frac{1}{2}mv^2$	2 -	$q \vec{E} = \frac{1}{2}mv^2$
3 -	$q\Delta V = qv \vec{B} $	4 -	$q \vec{E} = qv \vec{B} $
5 -	$qv \vec{B} = \frac{mv^2}{r}$	6 -	$q \vec{E} = \frac{mv^2}{r}$

Q2: Which equation configurations would be used for each of the various stages?

Acceleration
Plates


Velocity
Selector

Detection
Chamber

(Record your **three digit** answer in the Numerical Response boxes below)

1	4	5	
----------	----------	----------	--

Use the following information to answer Q3:



Q3: Using the information above, assign the model to the person that proposed the model:

Bohr

Dalton

Rutherford

J.J. Thomson

(Record your **four digit** answer in the Numerical Response boxes below)

4	1	3	2
----------	----------	----------	----------