

First Name: \_\_\_\_\_

Last Name: \_\_\_\_\_

**Worksheet - Unit 1 (Momentum) Basics**

**/12 marks**

**Q1:** A 1247kg Honda Civic DX Sedan is travelling at 100 km/h. What is its momentum?

- a.  $1.25 \times 10^4 \text{ kg}\cdot\text{m/s}$
- b.**  $3.46 \times 10^4 \text{ kg}\cdot\text{m/s}$
- c.  $1.25 \times 10^5 \text{ kg}\cdot\text{m/s}$
- d.  $4.49 \times 10^5 \text{ kg}\cdot\text{m/s}$

$$\frac{100 \text{ km}}{\text{h}} \times \frac{1 \text{ h}}{3600 \text{ s}} \times \frac{1000 \text{ m}}{1 \text{ km}} = 27.7 \text{ m/s}$$

$$\begin{aligned} p &= mv \\ &= (1247)(27.7) \\ &= 34,638.8 \text{ kg}\cdot\text{m/s} \\ &\approx 3.46 \times 10^4 \text{ kg}\cdot\text{m/s} \end{aligned}$$


**Q2:** A boy and his bicycle has a combined mass of 50.0kg and are travelling at 8.50 m/s. Upon applying the brakes, it takes the boy 3.25 seconds to come to a complete stop. What is the force of friction acting on the bicycle, in Newtons?

(Record your **three digit** answer in the Numerical Response boxes below)

1	3	1	
---	---	---	--

$$\begin{aligned} F_{\text{at}} &= m\Delta v \\ F(3.25) &= (50)(8.50) \\ F &= 130.7692.. \text{ N} \\ &\approx 131 \text{ N} \end{aligned}$$

Use the following information to answer Q3:



Q3: The final velocity of the 50g object is \_\_\_\_ m/s.

(Record your **three digit** answer in the Numerical Response boxes below)

1.75

$$F \Delta t = m \Delta v$$

$$0.0875 = (50 \times 10^{-3}) \Delta v$$


$$\Delta v = 1.75 \text{ m/s}$$

$$\Delta v = v_f - v_i$$

$$1.75 = v_f - (0)$$

$$v_f = 1.75 \text{ m/s}$$

Use the following information to answer Q4:



Q4: The final velocity of Object B is \_\_\_\_ m/s [R].

(Record your **three digit** answer in the Numerical Response boxes below)

1.87

$$p_i = p_f$$

$$(+6) + (-4.5) = (-4.1) + p_f$$


$$p_f = +5.6$$

$$v_f = +1.866 \text{ m/s}$$

$$\approx +1.87 \text{ m/s}$$

Key

Use the following information to answer Q5:




Q5: Which of the following statements is correct?

- a. Momentum is conserved and the collision is elastic.
- b. Momentum is conserved and the collision is not elastic.
- c. Momentum is not conserved and the collision is elastic.
- d. Momentum is not conserved and the collision is not elastic.

Q6: A 50.0kg skateboarder is travelling at 5.00 m/s when he picks up a stationary 4.50kg backpack. What is his final velocity?

(Record your **three digit** answer in the Numerical Response boxes below)

4 . 5 9




$$p_i = p_f$$

$$(+250) + (0) = (54.5)(v_f)$$

$$v_f = 4.59 \text{ m/s}$$

Use the following information to answer Q7:



Q7: What is the mass of Object B, in kg?

(Record your three digit answer in the Numerical Response boxes below)

1 . 2 5

$$p_i = p_f$$


$$(+10) + (-4m) = (3.08 + 1.54m)$$

$$10 - 4m = 3.08 + 1.54m$$

$$6.92 = 5.54m$$

$$m = 1.25 \text{ kg}$$


Use the following information to answer Q8:



Q8: The combined object moves at  $a.b$  m/s [ $ef^\circ$  N of E], where the values of  $a$ ,  $b$ ,  $e$ , and  $f$  are \_\_, \_\_, \_\_, and \_\_.


(Record your four digit answer in the Numerical Response boxes below)

3 1 5 0



$$\vec{v}_f = 3.1 \text{ m/s } [50^\circ \text{ N of E}]$$

Use the following information to answer Q9:



Q9: Object A moves at  $a.b \text{ m/s}$  [ $ef^\circ$  N of E], where the values of  $a$ ,  $b$ ,  $e$ , and  $f$  are \_\_, \_\_, \_\_, and \_\_.

(Record your **four digit** answer in the Numerical Response boxes below)

5	1	3	9
---	---	---	---

$$\vec{v}_f = 5.1 \text{ m/s } [39^\circ \text{ N of E}]$$


Key

Attention



Going to use "g m/s" instead of "kg m/s"


Use the following information to answer Q10:



Q10: An initially stationary object explodes. Object C moves at  $a.b$  m/s [ $ef^\circ$ ], where the values of  $a$ ,  $b$ ,  $e$ , and  $f$  are \_\_, \_\_, \_\_, and \_\_.


(Record your four digit answer in the Numerical Response boxes below)

1 9 4 1



$$\vec{v}_C = 1.9 \text{ m/s } [41^\circ]$$

Use the following information to answer Q11-Q12:



**Q11:** The ballistic pendulum rises to a height of  $a.bc \times 10^d \text{ m}$ , where  $a$ ,  $b$ ,  $c$ , and  $d$  are \_\_, \_\_, \_\_, and \_\_.

(Record your **four digit** answer in the Numerical Response boxes below)

4 1 0 2

COLLISION

$$p_i = p_f$$

$$(8.0 \times 10^{-3})(6.50) + (50 \times 10^{-3})(0) = (58 \times 10^{-3})v_f$$

$$v_f = 0.896551724138 \text{ m/s}$$

CONSERVATION OF E

$$E_k \rightarrow E_p$$

$$\frac{1}{2}mv^2 \rightarrow mgh$$

$$\frac{1}{2}(0.89655\dots)^2 \rightarrow (9.81)h$$

$$h = 4.096865\dots \times 10^{-2}$$

$$h \approx 4.10 \times 10^{-2} \text{ m}$$

**Q12:** What two **Physics Principles** are used to calculate the answer to **Q11**, listed in the order you used them?

(Record your **two digit** answer in the Numerical Response boxes below)

4 5