

First Name: _____

Last Name: _____

L15 - EQ - Photoelectric Effect 1

Q1: In the photoelectric effect the kinetic energy of emitted electrons is determined by the freq. of the incident light, and the number of emitted electrons is determined by the intensity of the incident light.

	Kinetic energy of emitted electrons is determined by:	Number of emitted electrons is determined by:
A.	Frequency	Frequency
B.	Frequency	Intensity
C.	Intensity	Frequency
D.	Intensity	Intensity

↓
Number of photons where each one "knocks off" an electron.

Q2: An incident photon of wavelength 450nm is incident on a Sodium surface with a work function of 2.28eV. Determine the kinetic energy of the emitted electrons, measured in electronvolts.

(Record you **three-digit** answer in the numerical response boxes below.)

0	.	4	8
---	---	---	---

$$E_{\text{photon}} = W + E_{\text{electron}}$$


$$\frac{hc}{\lambda} = W + E_k$$

$$\frac{(4.14 \times 10^{-15} \text{ eVs})(3 \times 10^8 \text{ m/s})}{(450 \times 10^{-9} \text{ m})} = 2.28 \text{ eV} + E_k$$

$$2.76 = 2.28 + E_k$$

$$E_k = 0.48 \text{ eV}$$

Use the following information to answer Q3 and Q4:



Q3: The slope of this graph is $a.b \times 10^{cd}$ Js, where a , b , c , and d are ____, ____, ____, and ____.

(Record your **four-digit** answer in the numerical response boxes below.)

6	6	3	4
---	---	---	---

Slope = Planck's constant
 $m = 6.63 \times 10^{-34}$ Js
 $m \approx 6.6 \times 10^{-34}$

Q4: The threshold frequency of the incident photons required for electron emission from a zinc target is $a.b \times 10^{cd}$ Hz, where a , b , c , and d are ____, ____, ____, and ____.

(Record your **four-digit** answer in the numerical response boxes below.)

1	0	1	5
---	---	---	---

$W = hf_0$
 $4.30 \text{ eV} = (4.14 \times 10^{-15} \text{ eVs}) f_0$
 $f_0 = 1.0386... \times 10^{15} \text{ Hz}$
 $f_0 = 1.0 \times 10^{15} \text{ Hz}$

MEY

Low τ large λ High τ Small λ
 RMI V U X G
 ↓
 Roy G. BIV
 750nm 400nm

Q5: The energy of a single red photon is

- a. 2.65×10^{-28} J
- b. 4.97×10^{-28} J
- c. 2.65×10^{-19} J
- d. 4.97×10^{-19} J

$$E = \frac{hc}{\lambda}$$

$$E = \frac{(6.63 \times 10^{-34})(3.0 \times 10^8)}{(750 \times 10^{-9})}$$

$$E = 2.652 \times 10^{-19} \text{ J}$$

Use the following information to answer Q6:

White light is shown through a double-slit apparatus (Diagram #1) and a glass prism (Diagram #2).

Diagram #1

... but Diffraction is Different

Diagram #2

Blue bends better...

In both cases, a rainbow spectrum appears.

Q6: The locations of **blue** and **red** light can be identified using the numbers given above.

Direction: <u>1</u>	<u>2</u>	<u>4</u>	<u>3</u>
Force: Location of blue light in Diagram #1	Location of red light in Diagram #1	Location of blue light in Diagram #2	Location of red light in Diagram #2

(Record your four digit answer in the Numerical Response boxes below)

1	2	4	3
---	---	---	---

MARKING:

Beginning	0.0 – 2.5
Progressing	3.0 – 4.0
Competent	4.5 – 5.5
Exemplary	6.0