


## L17 - Millikan's Oil-Drop Experiment

**Purpose:** Explain Millikan's oil-drop experiment and its significance relative to charge quantization.

**Key Result:**  
Charge is Quantized,  $e = 1.60 \times 10^{-19} \text{ C}$



*Indoor Skydiving Analogy regarding Unbalanced Forces*





*Descending*



*Not Accelerating*



# Millikan's Oil-Drop Experiment




# Millikan - With the Maths

$$V = \frac{4}{3}\pi r^3$$

$$\vec{E} = \frac{\vec{F}_e}{q}$$

$$\vec{g} = \frac{\vec{F}_g}{m}$$



$$\vec{F}_e = q\vec{E}$$


$$\vec{F}_g = m\vec{g}$$

Where  $|E| = \frac{\Delta V}{\Delta d}$

Where  $\rho = \frac{m}{V}$

Density =  $\frac{\text{Mass}}{\text{Volume}}$       $\rho = \frac{m}{V}$

Q1: An oil droplet of mass  $8.2 \times 10^{-15}$  kg is suspended in an electric field of  $1.0 \times 10^5$  N/C down. How many electrons has the oil drop gained or lost?



$$F_e = F_g$$

$$q|\vec{E}| = mg$$

$$q(1.0 \times 10^5) = (8.2 \times 10^{-15})(9.81)$$


$$|q| = 8.0442 \times 10^{-19} \text{ C} \quad |e| = 1.60 \times 10^{-19} \text{ C}$$

$$q = -5.027625e$$

$q = -5e$  Gained 5 electrons

3-digit  
5.00

Q2: A tiny plastic sphere of mass  $8.2 \times 10^{-15}$  kg is placed in an electric field of  $1.0 \times 10^5$  N/C [down] within a vacuum chamber. The sphere has 10 excess electrons. Determine whether the sphere will accelerate, and if so, in which direction.



$$F_g = mg$$

$$= (8.2 \times 10^{-15})(9.81)$$

$$= 8.0442 \times 10^{-14} \text{ N}$$

$$F_e = q|\vec{E}|$$

$$= (10 \times 1.60 \times 10^{-19})(1.0 \times 10^5)$$

$$= 1.60 \times 10^{-13} \text{ N}$$

Yes. Will accelerate upward.

**Q3:** During a Millikan oil – drop experiment, a student records the weights of five different oil drops. The student also records the electric field intensity necessary to hold each drop stationary between the two horizontal, parallel plates. Use a graph to calculate the elementary charge.


Weight (x 10 <sup>-14</sup> N)	Electric Field (x 10 <sup>5</sup> N/C)
1.7	1.1
5.6	3.5
6.1	3.8
2.9	1.8
4.0	2.5

Which variable **depends** on the other?

E-Field depends on Weight.  
Dependent (y-axis)                      Independent (x-axis)

$\vec{F}_e = \vec{F}_g$  ← x-axis

$q\vec{E} = m\vec{g}$  ← y-axis



✓

$F_e = F_g$

$qE = F_g$

$|E| = \frac{1}{q}(F_g) + 0$

$y = m(x) + b$

Slope =  $\frac{1}{q}$

$\frac{2.5 \times 10^5}{4.0 \times 10^{-14}} = \frac{1}{q}$

$6.25 \times 10^{18} = \frac{1}{q}$

$q = 1.60 \times 10^{-19} \text{ C}$